arXiv:astro-ph/0503044v2 8 Feb 2006

Effect of entrainment on stress and pulsar glitches in stratified neutron star crust

Nicolas Chamel$^{\star\,\dagger}$**, Brandon Carter***

* LuTh, Observatoire de Paris, 92195 Meudon, France
† Centrum Astronomiczne im M. Kopernika, Warszawa, Polska

Extended version, 10 December 2005

Abstract

The build up of the stress whose relaxation is presumed to account for pulsar frequency glitches can be attributed to various mechanisms, of which the most efficient involve differential rotation of the neutron superfluid in the inner layers of the (magnetically braked) solid crust of a rotating neutron star. In such a case it is usually supposed that the stress is attributable to pinning of superfluid vortices to crust nuclei, but it has been suggested that, even if the pinning effect is too weak, a comparably large stress might still arise just from the deficit of centrifugal buoyancy in the slowed down crust. The present work is a re-examination that investigates the way such processes may be affected by considerations that were overlooked in the previous work – notably uncertainties about the "effective" masses that have to be attributed to the "free" superfluid neutrons to allow for their entrainment by the ionic crust material. Though restricted to a Newtonian formulation, this analysis distinguishes more carefully than has been usual between true velocities, which are contravariantly vectorial, and so called "superfluid velocities" that are proportional to momenta, which are essentially covectorial, a technicality that is important when more than one independent current is involved. The results include a Proudman type theorem to the effect that the superfluid angular velocity must be constant on slightly deformed Taylor cylinders in the force free case, and it is shown how to construct a pair of integral constants of the motion that determine the solution for the pinned case assuming beta equilibrium.

1 Introduction

This work is intended as a comparison of the qualitative mathematical consequences of hypotheses of various alternative kinds (particularly concerning vortex pinning and chemical equilibrium) that have been invoked in the construction of differentially rotating neutron star models of the type developed to acccount for phenomena observed in pulsars.

It is hoped that this qualitative study will be helpful for the preparation of quantitative models designed to improve on those of the simple kind constructed by Prix et al [39], for which (as for the analogously constructed models [40] in a relativistic framework) separately conserved proton and superfluid neutron currents were able to deviate from Proudman - Taylor type corotation illustrated on figure 1, without giving rise to the (glitch producing) stresses that are actually anticipated. In earlier work [38] such stresses were obtained by including allowance for the effect of vortex pinning, but it has since been remarked [16] that deviations from corotation would give rise to stresses in any case (with or without pinning) when allowance is made for the consideration that the proton and neutron numbers will not be separately conserved but will undergo transfusion (by beta processes) so as to achieve chemical equilibrium.

The work of Prix [38] and of Carter et al [16] used a simplified treatment that neglected the mechanism of relative entrainment between the differentially rotating constituents, an effect which had already been considered in earlier work (both in a Newtonian framework [34] and also [30] in a relativistic treatment) but which had been considered to be only of secondary relevance as a minor quantitative correction. The effect of entrainment (but not transfusion) in the liquid core was allowed for in the more recent work of Comer & Andersson [18] and of Prix et al. [39], and it has since been recognised [17] to be particularly important in the crust layers that are the primary concern of the present analysis, which will consider the (glitch producing) stresses that can be expected to arise from the combined effect of entrainment, pinning, transfusion, and also nuclear stratification.

Explanations of the glitches of the pulsar period $P = 2\pi/\Omega$ observed in rotating neutron stars may be broadly classified into two main categories. In what may be described as the "deformation" category, the discontinuous spin up $\delta\Omega$ is attributed to a sudden change of the geometric configuration of the matter distribution [23], whereby the relevant moment of inertia is decreased so that – for a given angular momentum – the angular velocity Ω must increase. While conceivably sufficient to account for cases such as the glitches of order $\delta\Omega/\Omega \simeq 10^{-8}$ observed in the Crab pulsar, such a deformation mechanism is inadequate for explaining the frequent occurrence of the larger gliches $\delta\Omega/\Omega \simeq 10^{-6}$ observed in Vela like pulsars, not to mention the enormous glitch $\delta\Omega/\Omega \simeq 1.6 \times 10^{-5}$ recently observed in PSR J1806-2125 by Hobbs et al. [24].

To account for the frequent occurence of such large glitches, it has long been generally recognised [8] to be necessary to invoke a mechanism belonging to what may be referred to as the "transfer" category, involving interaction between two dynamically distinct constituents of the star. In such a case the discontinuous increase in the observed angular velocity Ω is attributed to transfer of angular momentum from a more rapidly rotating internal constituent to a more slowly rotating constituent that is directly coupled to the outer magnetosphere whose pulsations are directly observed. However it is not so clear what actually happens during the discontinous process involved. It has been suggested [42] that the stress due to pinning of neutron superfluid vortices to the ionic crust material may build up elastic stress to the point where the solid stucture itself breaks down (as in a terrestrial earthquake). In another much studied scenario [2, 37] it is postulated that before this can happen there will be a crisis of a different kind whereby a threshold is reached at which the vortices in the crust become collectively unpinned (whereas in the fluid core it is expected that vortices in the core will be subject to a strong dissipative drag, which will simulates pinning but will not be limited by any threshold). Another possibility [27, 26, 21, 22] is that pinning in the crust may be too weak to be effective, but it has been pointed out [16] that differential rotation might in principle by itself engender sufficient elastic stress to cause a breakdown of the solid structure even in the absence of pinning.

The main purpose of the present work is not to draw any specific conclusions about the relative plausibility of these various postulates in particular cases, but to consider, generally, how such alternatives will affect the predicted outcome, and how estimates of relevant quantities such as subcomponent moments inertia and stresses may need to be modified to take account of some effects that tended to be neglected in preceding discussions, including stratification and particularly the entrainment between the ionic crust material and the "free" superfluid neutrons, which thereby aquire "effective masses" that can be defined in various ways, but that in any case are likely[13, 14, 15] to be large enough to make an important difference.

In order to carry out such a revision, and as a secondary purpose in its own right, this work employs and develops a Newtonian formulation of the covariant kind [11, 12] whose use would be taken for granted in a relativistic treatment, in which care is taken to distinguish between true velocities, which are contravariantly vectorial, and so called "superfluid velocities" that are proportional to momenta, which are essentially covectorial.

Although deeply rooted in the well known principles of Lagrangian and Hamiltonian mechanics, the fundamental distinction between velocities and their canonical conjugates, namely momenta, has commonly been obscured in the context of superfluidity as traditionally formulated [34] in a Newtonian framework, for which it has long been customary to use the term "velocity" indiscriminately for anything having the physical dimensions thereoff, regardless of its actual mathematical role. The distinction is relatively unimportant for a single constituent perfect fluid or superfluid such as that of ordinary Helium at zero temperature, or even for a two contituent fluid in which entrainment is absent [16], but it becomes important when there are two or more independently moving fluids with entrainment, which means that the momenta need no longer be aligned with their conjugate velocities.

As well as simplifying the algebra in the work presented below, the introduction of the canonically covariant formulation also has the advantage of providing a helpful step towards the development of a fully General relativistic treatment, such as will ultimately be necessary for accuracy. Relativistic effects will be particularly important for the core, in which however, compared with the crust, the effect of entrainment is expected to be much more moderate [44, 19] at least in the outer part for which the basic physics is reasonably well understood, unlike the inner core for which many exotic possibilities have been envisages, such that of hyperon crystallisation [36], or of a LOFF phase in a color superconducting quark condensate [1].



Figure 1: Intersection of rotating neutron star model with a Taylor cylinder, over which the superfluid angular velocity must be uniform when the conditions for the Proudman theorem are applicable.

2 Two fluid model for stratified neutron star crust

2.1 The Lagrangian master function

The previous treatment referred to above [16] used a non relativistic Newtonian description in terms of free neutron and confined baryon 4-currents $n_{\rm f}^{\,\nu}$ and $n_{\rm c}^{\,\nu}$ respectively representing a neutron superfluid, and a normal crust component consisting of ionic nuclei characterised by a confined baryon number $A_{\rm c}$ and a positive charge number Z (in a degenerate electron background) that were treated as decoupled but interpenetrating barotropic perfect fluids interacting only via vortex pinning and long range gravitational forces, with the solid rigidity of the crust component taken into account by the inclusion of an extra stress force density.

One of the improvements introduced here is to drop the barotropy restriction for the ionic constituent by allowing its behavior to depend on the nuclear mass number $A_{\rm c}$ (which will tend to be larger in the deeper layers) thus allowing for the possibility of a stratification effect that may be important for stabilisation.

Another improvement that would be needed for a more accurate analysis is of course the use of a general relativistic treatment, of the kind that has long been commonly used with a simple perfect fluid description, and that has recently been developed for use with a two constituent fluid description of the kind needed fore treating the stellar core [30, 5]. However for the semilocal modelisation needed for the treatment of differential rotation in the crust layers with which we shall be concerned here, it would appear that corrections from general relativity will be much less important (at most a few tens of per cent) than those due to the entrainment. A recent program of work [13, 14, 15] (using a microphysical analysis combining methods from nuclear and solid state physics) on the relative motion of unbound neutrons in the inner crust layers (between the neutron drip density about 10^{11} g/cm^3 and nuclear saturation density about 10^{14} g/cm^3) has provided results [17] indicating that the effect of entrainment is likely to be very much stronger than had previously been supposed.

As in the preceding work [16], we shall therefore proceed here within a non relativistic Newtonian treatment, but instead of using a decoupled perfect fluid description we shall now include allowance for the effect of entrainment, using a non-dissipative two constituent description of the generic kind already developed for use in the liquid core [34, 18, 39] where the entrainment corrections are relatively moderate, though still potentially important as a cause for two-stream instability [6] if the relative velocity becomes too large.

For the reasons outlined in the introduction, it is both instructive and convenient (even though our treatment is non-relativistic) to use a canonical formalism of the 4-dimensionally covariant kind recently developed [11, 12] for the purpose of facilitating the exploitation of the variational formulation that is applicable to the non-dissipative limit. In this limit there is no resistance to relative motion, so that the only direct coupling between the two constituents is via the entrainment effect [7], whose microscopic origin is the Bragg scattering of dripped neutrons by crustal nuclei [13] which is the analog in the nuclear context of conduction electron scattering in ordinary solids.

It is also convenient to use a chemical constituent label ${\rm X}$ that runs over two values, namely ${\rm X}={\rm f}$ for the superfluid constituent and ${\rm X}={\rm c}$ for the crust constituent. Each constituent will have a baryon current density 4-vector having the form

$$n_{\rm X}^{\,\nu}=n_{\rm X}u_{\rm X}^{\,\nu} \tag{1}$$

in which the scalars $n_{\rm X}$ are the respective number densities of free baryons (only neutrons) and of baryons that are effectively confined within the atomic nuclei (namely all the protons, together with a roughly equal but somewhat larger number of confined neutrons, whose precise specification needs further clarification as discussed below) while the vectors $u_{\rm X}^{\,\nu}$ are the corresponding 4-velocities, whose components are labelled by a Greek index ν running over the values $0,1,2,3$. The conservation of the total baryon number current

$$n_{\rm B}^{\,\nu}=n_{\rm f}^{\,\nu}+n_{\rm c}^{\,\nu}\,, \tag{2}$$

is concisely expressible in this 4-dimensional notation as

$$\nabla_\nu n_{\rm B}^{\,\nu}=0\,. \tag{3}$$

In Aristotelian coordinates (representing the usual kind of 3+1 space time decomposition with respect to the rest frame of the star) the 4-velocity components will have the form $u_{\rm x}^{\,0} = 1$, $u_{\rm x}^{\,i} = v_{\rm x}^{\,i}$ where the $v_{\rm x}^{\,i}$ are ordinary 3-velocity components, using a Latin index $i = 1, 2, 3$. This means that time component of the current density is simply equal to the corresponding particle number density $n_{\rm x}^{\,0} = n_{\rm x}$, while the space components are the those of the current density 3-vector $n_{\rm x}^{\,i}$.

In addition to the six independent vectorial components provided by the space vectors $n_{\rm c}^{\,i}$ and $n_{\rm f}^{\,i}$ and the pair of scalars provided by the "confined" and "free" baryon number densities $n_{\rm c}$ and $n_{\rm f}$, our model will contain a nineth independent variable, namely the scalar provided by the nuclear mass number $A_{\rm c}$, or equivalently by the number density $n_{\rm I}$ of nuclei, whose number current will be given by

$$n_{\rm I}^{\,\nu} = n_{\rm I} u_{\rm c}^{\,\nu} \,, \qquad\qquad n_{\rm I} = n_{\rm c}/A_{\rm c} \,. \tag{4}$$

We shall take the mass per baryon to be the same (considering the electron mass and the proton neutron mass difference to be negligible compared with the ordinary proton mass) with fixed value $m^{\rm c} = m^{\rm f} = m$, so that the corresponding mass density contributions will simply be given by $\rho_{\rm x} = m\, n_{\rm x}$. The total mass density, as given explicitly by

$$\rho = m\, n_{\rm B} = \rho_{\rm f} + \rho_{\rm c} \,, \tag{5}$$

will be important as the Poissonian source of the Newtonian gravitational potential scalar, ϕ. Within this Newtonian background field, the total force balance equation will take the form

$$\nabla_\mu T^\mu{}_\nu = f_\nu - \rho \nabla_\nu \phi \,, \tag{6}$$

in which f_ν is the non gravitational external 4-force density – if any – that may be acting on the system, and $T^\mu{}_\nu$ is the stress momentum energy density tensor as constructed in terms of the 4-dimensional formalism recapitulated below.

The dynamics of system will be governed – in the manner described in our preceding work [11, 12] – by a Lagrangian master function

$$\Lambda = \Lambda_{\rm mat} - \phi\, \rho \,, \qquad\qquad \Lambda_{\rm mat} = \tilde\Lambda_{\rm kin} + \tilde\Lambda_{\rm int} \tag{7}$$

in which $\Lambda_{\rm kin}$ is the ordinary Galilean frame dependent kinetic energy density, namely

$$\tilde\Lambda_{\rm kin} = \tfrac{1}{2} \left(\rho_{\rm f} v_{\rm f}^{\,2} + \rho_{\rm c} v_{\rm c}^{\,2} \right) \,, \tag{8}$$

and $\tilde\Lambda_{\rm int}$ is some appropriately specified internal action density contribution that must be Galilean frame independent, which means that it can depend only on the scalar number densities $n_{\rm I}$, $n_{\rm f}$, $n_{\rm c}$, and on the velocity difference

$$[v^i] = v_{\rm f}^{\,i} - v_{\rm c}^{\,i} \tag{9}$$

so that in terms of an "entrainment density" function $\bar\rho_{\rm fc}$ and a set of chemical potentials $\tilde\chi^{\rm f}$, $\tilde\chi^{\rm c}$, $\tilde\chi^{\rm I}$ its generic variation will take the form

$$\delta\tilde\Lambda_{\rm int} = \bar\rho_{\rm fc}[v_i]\, \delta[v^i] - \tilde\chi^{\rm f} \delta n_{\rm f} - \tilde\chi^{\rm c} \delta n_{\rm c} - \tilde\chi^{\rm I} \delta n_{\rm I} \,, \tag{10}$$

which gives the corresponding formula

$$\delta\Lambda_{\rm mat} = n_{\rm f} \mu^{\rm f}_i \delta v_{\rm f}^{\,i} + n_{\rm c} \mu^{\rm c}_i \delta v_{\rm c}^{\,i} - \tilde\mu^{\rm f} \delta n_{\rm f} - \tilde\mu^{\rm c} \delta n_{\rm c} - \tilde\mu^{\rm I} \delta n_{\rm I} \,, \tag{11}$$

in which the (Galilean frame dependent) material energy coefficients are given in terms of the corresponding (frame independent) chemical potentials by

$$\tilde\mu^{\rm f} = \tilde\chi^{\rm f} - \tfrac{1}{2} m\, v_{\rm f}^{\,2} \,, \quad \tilde\mu^{\rm c} = \tilde\chi^{\rm c} - \tfrac{1}{2} m\, v_{\rm c}^{\,2} \,, \quad \tilde\mu^{\rm I} = \tilde\chi^{\rm I} \,, \tag{12}$$

and the associated space momenta are given by

$$\mu^{\rm f}_i = m\, v_{{\rm f}\, i} + \frac{\bar\rho_{\rm fc}}{n_{\rm f}} [v_i] \,, \qquad\qquad \mu^{\rm c}_i = m\, v_{{\rm c}\, i} - \frac{\bar\rho_{\rm fc}}{n_{\rm c}} [v_i] \,. \tag{13}$$

In the low relative velocity regimes that are relevant it will always be possible to take the internal energy contribution to have the quadratic form

$$\tilde\Lambda_{\rm int} = \tfrac{1}{2} \bar\rho_{\rm fc} [v]^2 - \tilde U_{\rm ins} \,, \tag{14}$$

with $\bar\rho_{\rm fc}$ and the static contribution $\tilde U_{\rm ins}$ specified by appropriate equations of state as functions only of the scalar densities $n_{\rm I}$, $n_{\rm c}$, $n_{\rm f}$. It is to be remarked that to translate our notation to that used by Prix et al. [39] we would need to make the conversions $\tilde\chi^{\rm f} \mapsto \mu^{\rm f}$, $\tilde\chi^{\rm c} \mapsto \mu^{\rm c}$ and $\tilde\Lambda_{\rm int} \mapsto -\mathcal{E}$ where $\mathcal{E}$ is what these authors have referred to rather loosely as "internal energy" density, although it is something that should not be confused with the true internal energy density $\tilde U_{\rm int}$, which will actually be given by

$$\tilde U_{\rm int} = \tfrac{1}{2} \bar\rho_{\rm fc} [v]^2 + U_{\rm ins} = \bar\rho_{\rm fc} [v]^2 - \Lambda_{\rm int} \,. \tag{15}$$

2.2 Currents and their conjugate momenta in 4 dimensions

Instead of treating quantities such as energy and momentum separately, as has traditionally been done in a Newtonian framework, it has been found to be technically advantageous, as remarked above, to unify them in a 4-dimensional treatment [11, 12] (which was originally inspired by the corresponding relativistic theory [30]). The first step in this 4 dimensional treatment is to rewrite the generic variation (11) in tems of 4-current variations in the form

$$d\Lambda_{\rm mat} = \mu^{\rm f}_\nu \, dn_{\rm f}^{\,\nu} + \mu^{\rm c}_\nu \, dn_{\rm c}^{\,\nu} - \tilde\mu^{\rm I} \, dn_{\rm I} \,, \tag{16}$$

which provides a direct specification of the material 4-momentum covectors $\mu^{\rm f}_{\,\nu}$ and $\mu^{\rm c}_{\,\nu}$ that play a crucial role in the concise canonical formalism to be described below. In terms of the quantities introduced above their time components will be given by

$$\mu^{\rm f}_{\ 0} = -(\tilde\mu^{\rm f} + \mu^{\rm f}_{\,i} v_{\rm f}^{\,i}) \, , \qquad \mu^{\rm c}_{\ 0} = -(\tilde\mu^{\rm c} + \mu^{\rm c}_{\,i} v_{\rm c}^{\,i}) \, , \tag{17}$$

while their space components will simply be given by (13).

These material 4-momenta, $\mu^{\rm f}_{\,\nu}$ and $\mu^{\rm c}_{\,\nu}$, give rise to corresponding complete "free" and "confined" baryon momenta, $\pi^{\rm f}_{\,\nu}$ and $\pi^{\rm c}_{\,\nu}$, whose time components include allowance for gravitational energy according to the specifications

$$\pi^{\rm f}_{\ 0} = \mu^{\rm f}_{\ 0} - m\phi \, , \qquad \pi^{\rm c}_{\ 0} = \mu^{\rm c}_{\ 0} - m\phi \, , \tag{18}$$

while their space components are given simply by

$$\pi^{\rm f}_{\ i} = \mu^{\rm f}_{\ i} \, , \qquad \pi^{\rm c}_{\ i} = \mu^{\rm c}_{\ i} \, . \tag{19}$$

It is also useful to introduce the total ionic momentum covector $\tilde\pi^{\rm I}_{\ \nu}$, which is defined so as to include allowance for nuclear energy by

$$\tilde\pi^{\rm I}_{\ 0} = A_{\rm c} \pi^{\rm c}_{\ 0} - \tilde\mu^{\rm I} \, , \qquad \tilde\pi^{\rm I}_{i} = A_{\rm c} \pi^{\rm c}_{\ i} \, . \tag{20}$$

This enables us to express the complete stress energy tensor of the system in the very concise form

$$T^\mu_{\ \nu} = n_{\rm f}^\mu \pi^{\rm f}_{\ \nu} + n_{\rm I}^\mu \tilde\pi^{\rm I}_{\ \nu} + \Psi\, \delta^\mu_\nu \, , \tag{21}$$

in which the material pressure function is given by

$$\Psi = \Lambda_{\rm mat} - n_{\rm f}^{\,\nu} \mu^{\rm f}_{\,\nu} - n_{\rm c}^{\,\nu} \mu^{\rm c}_{\,\nu} + n_{\rm I}\, \tilde\mu^{\rm I} \, . \tag{22}$$

It follows that the force balance equation (6) can be rewritten in the form

$$f^{\rm f}_{\ \nu} + f^{\rm I}_{\ \nu} = f_\nu \, , \tag{23}$$

in which the 4-force density $f^{\rm f}_{\,\nu}$ acing on the"free" neutrons is specified in terms of their vorticity tensor $\varpi^{\rm f}_{\ \mu\nu}$ by

$$f^{\rm f}_{\ \nu} = n_{\rm f}^\mu \varpi^{\rm f}_{\ \mu\nu} + \pi^{\rm f}_{\ \nu} \nabla_\mu n_{\rm f}^\mu \, , \qquad \varpi^{\rm f}_{\ \mu\nu} = 2\nabla_{[\mu} \pi^{\rm f}_{\ \nu]} \, , \tag{24}$$

while the combined 4-force density acting on the ions and their "confined" baryons will be specified by the formula

$$f^{\rm I}_{\ \nu} = 2 n_{\rm I}^\mu \nabla_{[\mu} \tilde\pi^{\rm I}_{\ \nu]} + n_{\rm I}^\mu \pi^{\rm c}_{\ \mu} \nabla_\nu A_{\rm c} \, . \tag{25}$$

This can be decomposed in the form

$$f^{\rm I}_{\ \nu} = f^{\rm c}_{\ \nu} + n_{\rm I} \nabla_\nu \tilde\mu^{\rm I} - \delta^0_\nu \nabla_\mu (\tilde\mu^{\rm I} n_{\rm I}^{\,\mu}) \, , \tag{26}$$

in which the last term contributes only to the time component, while the only part that remains in the absence of stratification is the contribution of first term, which is given by an expression of the standard form

$$f^{\rm c}_{\ \nu} = 2 n_{\rm c}^\mu \nabla_{[\mu} \pi^{\rm c}_{\ \nu]} + \pi^{\rm c}_{\ \nu} \nabla_\mu n_{\rm c}^\mu \, . \tag{27}$$

2.3 Equations of state and effective mass

To complete the specification of the variables introduced in the previous section it is evidently necessary to prescribe the particular form of the material action density $\Lambda_{\rm mat}$, which is – for this purpose – most conveniently decomposible in the form

$$\Lambda_{\rm mat} = U_{\rm dyn} - \tilde U_{\rm ins} \, , \tag{28}$$

wherein $\tilde U_{\rm ins}$ is the static internal energy contribution introduced in (14), which is specified by an appropriate "primary" equation of state as a function of the three relevant number densities $n_{\rm f}$, $n_{\rm c}$ $n_{\rm I}$, while the remaining dynamical energy contribution is given [15] in terms of the "normal" (crust) velocity 3 vector $v_{\rm c}^i$ and the relative current 3 vector

$$n^i = n_{\rm f} [v^i] \, , \tag{29}$$

by the formula

$$U_{\rm dyn} = \rho\, v_{{\rm c}\, i} v_{\rm c}^i/2 + m\, n_i v_{\rm c}^i + n_i n^i/2\mathcal{K} \, , \tag{30}$$

in which the $\mathcal{K}$ is the "mobility coefficient". This quantity $\mathcal{K}$ is given by a "secondary" equation of state as a function of the relevant number densities $n_{\rm I}$, $n_{\rm c}$, and $n_{\rm f}$.

In the analysis that follows a particularly important role will be played by the "free" neutron 3-momentum covector $\mu^{\rm f}_{\,i}$ for which, on a local "mesoscopic" scale (small compared with the intervortex separation) the superfluidity property entails the irrotationality condition $\nabla_{[i} \mu^{\rm f}_{\ j]} = 0$. It follows from (30) that the contravariant version (as obtained by contracting with the ordinary Euclidean space metric γ^{ij}) of this (superfluid) momentum will be given by an expression of the form

$$\mu^{{\rm f}\, i} = m_\star v_{\rm f}^i + (m - m_\star) v_{\rm c}^i \tag{31}$$

in terms of an effective mass variable,

$$m_\star = n_{\rm f}/\mathcal{K} \, , \tag{32}$$

whose difference from the ordinary baryon mass m is proportional to the "entrainment density"

$$\bar\rho_{\rm fc} = \rho_{\rm f} (m_\star - m)/m \, , \tag{33}$$

which – in much of the inner crust – now seems likely [17] to be much larger than had previously been supposed.

The reason why it is convenient for the formulation of the equations of state to start with the specification of $\mathcal{K}$ rather than the effective mass $m_\star$ or the corresponding "entrainment density" $\bar\rho_{\rm fc}$ in the decomposition (14) is that – unlike $m_\star$ and $\bar\rho_{\rm fc}$ – the "mobility coefficient" $\mathcal{K}$ has the advantage of being physically well defined in a manner that does not depend on the choice of the chemical basis that will be discussed in Section 2.6.

Another kind of decomposition that is also independent of the choice of chemical basis, and that is instructive for the purpose of comparison with the notation used in related work, including the seminal article of [7], is to introduce the concept of the so called "superfluid 3-velocity" $V^{\rm S}_i$, which in stricter terminology should be referred to as "superfluid momentum per unit mass", as defined by setting

$$\mu^{\rm f}_{\,i} = m\, V^{\rm S}_i \qquad\qquad V^{{\rm S}\,i} = v_{\rm c}^{\,i} + \frac{m_\star}{m}[v^i]\,. \tag{34}$$

This enables the dynamical energy density (30) to be rewritten (without a cross term) in the seductively simple form

$$U_{\rm dyn} = \frac{m}{2}\left(n_{\rm S} V^{\rm S}_i V^{{\rm S}\,i} + n_{\rm N} v_{{\rm c}\,i} v_{\rm c}^{\,i}\right)\,, \tag{35}$$

in which the so called "superfluid particle number density" $n_{\rm S}$ and the corresponding"normal particle number density" $n_{\rm N}$ are defined by

$$n_{\rm S} = m\mathcal{K} = \rho_{\rm f}/m_\star \qquad\qquad n_{\rm N} = n_{\rm B} - n_{\rm S}\,. \tag{36}$$

2.4 Dynamical Equations

Even after the primary (3 variable) equation of state function $\tilde{U}_{\rm ins}\{n_{\rm I}, n_{\rm c}, n_{\rm f}\}$ and the secondary (2 variable) equation of state function $\mathcal{K}\{n_{\rm I}, n_{\rm c}\}$ have been specified, the formulae given so far will merely provide a coherent set of definitions, but – except for the total baryon conservation law (3) – they contain no dynamical information. To govern the evolution of the nine independent variables (namely $n_{\rm I}$ and the components of the 4-vectors $n_{\rm f}^{\,\nu}$ and $n_{\rm c}^{\,\nu}$) of the system we still need eight more conditions, which can be provided by specification of the values of the 4-forces $f^{\rm f}_{\ \nu}$ and $f^{\rm c}_{\ \nu}$.

In the absence of an external force f_ν in (23) the sum of the contributions $f^{\rm f}_{\ \nu}$ and $f^{\rm c}_{\ \nu}$ would have to vanish, a condition that will automatically be satisfied if these contributions are specified just by the variational principle, according to which they should each vanish separately, $f^{\rm f}_{\ \nu} = f^{\rm c}_{\ \nu} = 0$.

In a dissipative model o the kind recently developed by [12] (such as might be needed to allow for the mutual resistivity that would be present if the temperature were too high for superconductivity) the contributions $f^{\rm f}_{\ \nu}$ and $f^{\rm c}_{\ \nu}$ could still add up to zero even though they would no longer vanish separately. However we wish to consider more general situations in which a non-vanishing total force density f_ν may be present for two reasons. One reason is to allow for the "secular" (very long timescale) effect of weak magnetic braking of pulsar rotation. However the main reason for introducing the external force contribution f_ν in the present analysis is to take account of the presence, due to the solidity of the crust, of additional stresses that we wish to estimate but for which a complete treatment would be beyond the scope of the purely fluid model used here.

In conjunction with (3), strict application of the variation principle would entail as consequence that each of the three currents $n_{\rm I}^{\,\nu}$, $n_{\rm f}^{\,\nu}$, $n_{\rm c}^{\,\nu}$ should be separately conserved. This would no doubt be a very good approximation for the purpose of treating high frequency oscillation modes. However, in the medium to long timescale processes we wish to consider here, whereas it will still be reasonable to postulate conservation of the number of ionic nuclei, meaning that we we shall have

$$\nabla_\nu n_{\rm I}^{\,\nu} = 0\,, \tag{37}$$

on the other hand it may be realistic to allow for the possibility that – due to beta processes whereby neutrons are transformed into neutrons or vice versa – the "free" and "confined" baryon currents may not be separately conserved, but will evolve in such a way as to diminish the magnitude of the relevant chemical affinity[12], $\tilde{\mathcal{A}}$ say, meaning the chemical potential difference between the two constituents, as measured in the relevant "normal" (non superfluid) rest frame, namely that of the ionic lattice with unit velocity 4-vector $u_{\rm c}^{\,\nu}$, so that it will be given by

$$\tilde{\mathcal{A}} = u_{\rm c}^{\,\nu}[\mu_\nu]\,, \qquad\qquad [\mu_\nu] = \mu^{\rm f}_{\ \nu} - \mu^{\rm c}_{\ \nu}\,. \tag{38}$$

As discussed in the treatment of the analogous relativistic model [30] (with $n_{\rm I}$ and $\mu^{\rm I}$ respectively replaced by entropy density and temperature) a non-dissipative model that is self contained – so that total force f_ν vanishes – will be obtainable in a manner that is compatible either with the short term conservation conditions $\nabla_\nu n_{\rm f}^{\,\nu} = \nabla_\nu n_{\rm c}^{\,\nu} = 0$, or alternatively with the long term equilibrium condition $\tilde{\mathcal{A}} = 0$ that is more relevant here, provided the three force acting on the superfluid neutron current is specified in such a way that

$$n_{\rm f}^{\,\mu}\varpi^{\rm f}_{\ \mu\nu} n_{\rm c}^{\,\nu} = 0\,. \tag{39}$$

2.5 Superfluidity condition

On a mesoscopic scale (large compared with the ionic spacing but small compared with the distance between quantised vortices) the superfluidity property of the "free" neutrons entails that their momentum covector $\pi^{\rm f}_{\ \nu}$ must be locally proportional to the gradient of a quantum phase scalar, and hence that their vorticity 2-form must vanish, $\varpi^{\rm f}_{\ \mu\nu} = 0$, which automatically ensures that the condition (39) will be satisfied.



Figure 2: Sketch of the 2-surface swept out by a quantised vortex line.

What we are concerned with here however is the macroscopic level – meaning lengthscales large compared with the intervortex separation – at which there will be a non-vanishing vorticity 2-form, and a corresponding spacelike vorticity vector

$$w^\lambda = \tfrac{1}{2}\varepsilon^{\lambda\mu\nu}\varpi^{\rm f}{}_{\mu\nu}\,, \tag{40}$$

whose components in Aristotelian coordinates are simply $w^0 = 0$ and $w^i = \varepsilon^{ijk}\nabla_j v_{{\rm f}k}$, representing the mean density of circulation around the quantised vortices. The superfluidity property does however entail an algebraic restriction to the effect that we must have

$$w^\mu \varpi^{\rm f}{}_{\mu\nu} = 0\,, \tag{41}$$

a condition need not be satisfied for a more general fluid motion (such as would be possible for the free neutrons in a very young neutron star that has not yet dropped below the temperature of the BCS pairing transition). This restriction expresses the requirement that, instead of having matrix rank 4 as in the normal case, the antisymmetric (so necessarily even ranked) vorticity tensor $\varpi^{\rm f}{}_{\mu\nu}$ should have its rank reduced to 2 in the superfluid case, a condition that is mathematically necessary for it to be tangential to a set of 2-surfaces which in this instance are swept out by the quantised vortex lines as illustrated on figure 2. This implies the existence of vorticity transport 4-vector field $u_{\rm w}^\nu$ subject to the usual time normalisation condition $u_{\rm w}^0 = 1$ characterised by the defining property

$$u_{\rm w}^\mu \varpi^{\rm f}{}_{\mu\nu} = 0 \tag{42}$$

which does not fix it completely but evidently allows a freedom of adjustment by addition of a spacelike vector field aligned with the direction of the vortex lines as given by w^ν.

The supposition that the vortices will be "pinned" in the sense of being transported with the ionic lattice is evidently expressible by adopting a dynamic equation of the form (42) with

$$u_{\rm w}^\mu = u_{\rm c}^\mu \tag{43}$$

which still automatically ensures that the condition (39) will be satisfied, but which will generically entail the involvement of a non-vanishing force density that will be attributable to the Magnus effect as discussed in the appendix. This condition (39) particularly plausible as a description of evolution on very long timescales, for which the beta equilibrium condition,

$$\tilde{\mathcal{A}} = 0\,, \tag{44}$$

will also be appropriate.

For evolution on the more moderate timescales that will be relevant for many astrophysical processes the situation is not so clear. An obvious – though not necessarily realistic – alternative way of satisfying (39) is to apply the strict variation principle to the effect that $f^{\rm f}_{\ \nu}$ should vanish, which instead of (43) as well as the separate "free" neutron conservation condition $\nabla_\nu n_{\rm f}^\nu = 0$ also provides a dynamical equation of the form (42) but with

$$u_{\rm w}^\mu = u_{\rm f}^\mu\,, \tag{45}$$

and which instead of (44) provides a separate conservation law

$$\nabla_\nu n_{\rm f}^\nu = 0\,. \tag{46}$$

However the realism of such an alternative is questionable on the basis of microphysical considerations which suggest that both the vortex slipping implied by (45) and the beta disequilibrium entailed by (46) would in practise give rise to dissipative processes that would require a more elaborate treatment. This issue is hinted at, on a purely macroscopic level, by the consideration that the actual physical meaning of the simple set of non dissipative evolution equations provided by (45) and (46) is open to question, due to its gauge dependence in the sense discussed in the next section.

2.6 Chemical basis dependence

A problem with the force free postulate, namely the use of (45) in conjunction with (46) is that (unlike the use of (43) in conjunction with (44) which will be appropriate for longer timescale processes) its physical meaning depends on how many of the neutrons are considered to be "free". To be more explicit, it depends on the choice of the dimensionless parameter $a_{\rm c}$ in the chemical base transformation

$$n_{\rm c}^\nu = a_{\rm c}\, n_{\rm p}{}^\nu \qquad n_{\rm f}^\nu = n_{\rm n}^\nu + (1-a_{\rm c}) n_{\rm p}{}^\nu\,, \tag{47}$$

whereby the "confined" and "free" baryon currents are specified in terms of the more obviously well defined chemical basis constituted by the proton current ${n_{\rm p}}^\nu$ and the total ("confined" plus "free") neutron current $n_{\rm n}^{\,\nu}$.

Obvious simple possibilities for the choice of such a "chemical gauge" include the "comprehensive" gauge, $a_{\rm c}$ in which all the neutrons are counted as "free", the "pairing gauge" $a_{\rm c} = 2$ in which the most strongly bound neutron states, namely those paired with proton states, are counted as "confined", but all the others are considered to be free. A more intuitively "natural" – but more fuzzily defined – possibility would be to reserve the qualification "free" for neutrons that are actually located outside the – somewhat blurred – boundaries of the nuclei that contain the protons. However what really matters is not location in ordinary space but in phase space: the "operational" criterion for a neutron state to be effectively "free" is that it should have sufficient energy to overcome the barriers separating the ionic potential wells either classically or by quantum tunnelling within the relevant timescale.

On this "operational" basis it is clear that all the neutrons will be effectively "free" in the fluid core and that none of them will be effectively "free" in the outer crust below the "drip" density threshold, while at densities just above this threshold there will be a clear cut critical energy above which the neutrons will be able to travel over ionic separation distances on a microscopically short timescale and below which their wave functions will be exponentially suppressed outside the ionic wells so that they will be able to tunnell only on cosmologically long timescales. However in the deeper and denser layers of the crust (which are likely to be particularly important for glitch processes) this "operational" distinction will no longer be so clear cut, because the ionic wells will get too close for the exponential suppression outside to be fully effective, so that there will be marginally bound states with intermediate penetration timescales that are macroscopically long but cosmologicaly short. This means that the most appropriate way of precisely defining an "operational" basis will depend on the timescales involved in the astrophysical context under consideration.

In view of the debatability of this question about which basis may be most appropriate, it is important to observe that the counting convention used to define the chemical gauge parameter $a_{\rm c}$ in (47) makes no difference at all to the specification of the "normal" velocity $v_{\rm c}^{\,i}$ and the relative current n^i defined by (29) and hence that it has no effect on any of the terms in the original specification (30) for the dynamical energy density. It is no less important to observe that uncertainties about the appropriate choice for $a_{\rm c}$ make no difference to the specification of the "free" momentum covector ${\mu^{\rm f}}_\nu$, nor in consequence, to the vorticity form ${\varpi^{\rm f}}_{\mu\nu}$ and the so called "superfluid 3-velocity" $V^{\rm s}_i$, which means that the quantities appearing in the rewritten formula (35) will also be unaffected.

The rule chosen for the specification of $a_{\rm c}$ and hence of $n_{\rm f}$ will evidently affect the individual terms in the formula (21) for the stress energy tensor, but it can be seen that the total, ${T^\mu}_\nu$, will be unaffected by the value of $a_{\rm c}$ so long as it is held fixed, and even (though this is less obvious) if it is allowed to vary, so long as this ratio,

$$a_{\rm c} = n_{\rm c}/n_{\rm p} = A_{\rm c} n_{\rm I}/n_{\rm p}\,, \tag{48}$$

is postulated to depend only on the nuclear charge number Z, which is itself definable as the ratio

$$Z = n_{\rm p}/n_{\rm I}\,. \tag{49}$$

It thus follows from (6) that provided $a_{\rm c}$ (or equivalently $A_{\rm c}$) is chosen as some function just of the single variable Z, although the choice of this function will indeed affect the specification of the separate force density contributions ${f^{\rm f}}_\nu$ and ${f^{\rm c}}_\nu$, nevertheless it will not have any effect on their total (23). This means that it will not have any effect on the complete set of evolution equations given by specifying the total external force f_ν (e.g. by postulating that it should vanish) in conjunction with the ionic number conservation law (37) together with the (long timescale) pinning and beta equilibrium conditions (43) and (44). However if (44) is replaced by (46), then the physical specification of the system would depend on the ansatz for the function $a_{\rm c}\{Z\}$, which for very short timescales could realistically be taken to be simply the "comprehensive" value $a_{\rm c} = 1$, corresponding just to separate conservation of protons and neutrons. Similarly if (43) is replaced by (45) the result would again depend on the ansatz for $a_{\rm c}\{Z\}$, but in this case the appropriate choice is a question needing further investigation at a microscopic level.

The strategy of the following work is not to adopt any particular choice of gauge but to see how far it is possible to draw general conclusions that will be valid whatever the choice of the function $a_{\rm c}\{Z\}$ and whatever the postulate adopted for the specification of the vortex drift velocity $u_{\rm w}^{\,\mu}$ in (42).

3 Axisymmetric configurations

3.1 Angular momentum convection and the lag formula

As in the preceding work [16] on this problem, we now restrict our attention to configurations that are axisymmetric in the sense of being invariant with respect to the action of a rotation symmetry generator ϱ^ν of the usual kind, meaning one that is spacelike, $\varrho^0 = 0$, with space



Figure 3: Axisymmetry generator ϱ^i, whose magnitude is equal to the cylindrical radius ϱ and the unit vector ν^i directed along the rotation axis.

components ϱ^i given in terms of the unit vector ν^i along the relevant symetry axis, with respect to Cartesian coordinates r^i centered on the axis, by an expression of the form $\varrho^i=\varepsilon^{ijk}\nu_j r_k$ as shown on figure 3. This means that this symmetry generator will have a scalar magnitude ϱ that can be interpreted as a cylindrical radial coordinate (such as was denoted by ϖ in the preceding work by Carter et al. [16]) so that it will be given in terms of the relevant angle of latitude θ by $\varrho=r\cos\theta$.

Invariance of a field with respect to such an action is expressible in any (Cartesian or other) coordinate system as the vanishing of its Lie derivative with respect to the generator. In the particular case of the superfluid 4-momentum covector $\pi^{\rm f}_{\ \nu}$ the axisymmetry condition will therefore be expressible as

$$\varrho^\nu\nabla_\nu\pi^{\rm f}_{\ \mu}+\pi^{\rm f}_{\ \nu}\nabla_\mu\varrho^\nu=0\,. \tag{50}$$

In terms of the angular momentum per free (superfluid) particle, as defined (independently of any choice of the chemical gauge parameter $a_{\rm c}$) by

$$\ell=\varrho^\nu\pi^{\rm f}_{\ \nu}\,, \tag{51}$$

(so that on a microscopic scale ℓ will be quantised as a half integer multiple of the Dirac-Planck constant $\hbar$) the axisymmetry condition (50) can evidently be rewritten as

$$\nabla_\mu\ell=\varpi^{\rm f}_{\ \mu\nu}\varrho^\nu\,. \tag{52}$$

We can use this in conjunction with (42) to obtain a generalised Bernoulli type [12] conservation law to the effect that this superfluid angular momentum scalar ℓ will be convected by (in the sense of remaining constant along) the vortex flux trajectories:

$$u_{\rm w}^{\ \nu}\nabla_\nu\ell=0\,. \tag{53}$$

We are concerned here with configurations that differ only very slightly from a state of circularity, meaning a state in which the 3-velocities are all aligned with the axisymmetry generator ϱ^i, so that we can write

$$v_{\rm x}^{\ i}=\Omega_{\rm x}\,\varrho^i+v_{{\rm x}\perp}^{\ \ i}\,,\qquad \varrho_i\,v_{{\rm x}\perp}^{\ \ i}=0\,, \tag{54}$$

where $\Omega_{\rm x}$ is the relevant angular velocity, and the residual non circular velocity contribution $v_{{\rm x}\perp}^{\ \ i}$ is supposed to be very small except perhaps in very brief intervals during a glitch. The non circular part, if any, will evidently make no contribution to the superfluid angular momentum scalar, which, according to (64) will be given simply by

$$\ell=\varrho^2\Big(m_\star\Omega_{\rm f}+(m-m_\star)\Omega_{\rm c}\Big)\,. \tag{55}$$

Since the axisymmetry evidently implies $\varrho^i\nabla_i\ell=0$, the conservation law (53) can be rewritten in terms of the vortex drift 3-velocity $v_{\rm w}^{\ i}$ as a slow variation rule of the form

$$\dot\ell=-v_{{\rm w}\perp}^{\ \ i}\nabla_i\ell\,, \tag{56}$$

with the usual convention that a dot denotes partial differentiation with respect to time at a fixed space position.

In the rotating neutron star configurations under consideration it will be justifiable to use a small perturbation approximation, not just for the non-circular velocity contributions, but also for deviations of the circular part from rigid rotation with a uniform fixed angular velocity value, $\bar\Omega$ say. This means that for each constituent we can write

$$\Omega_{\rm x}=\bar\Omega+\Delta\Omega_{\rm x} \tag{57}$$

where, like $v_{{\rm w}\perp}^{\ \ i}$ the deviation $\Delta\Omega_{\rm x}$ is to be considered as relatively small. It follows that to first perturbative order we can rewrite the slow variation law (56) as

$$\dot\ell=-m\bar\Omega\,v_{{\rm w}\perp}^{\ \ i}\nabla_i\varrho^2\,. \tag{58}$$

Using square brackets for differences between the constituents according to the convention

$$[\Omega]=\Omega_{\rm f}-\Omega_{\rm c}\,,\qquad [\dot\Omega]=\dot\Omega_{\rm f}-\dot\Omega_{\rm c}\,, \tag{59}$$

we can write (58) more explicitly as

$$m_\star[\dot\Omega]+m\dot\Omega_{\rm c}=-\frac{2m\bar\Omega}{\varrho}v_{{\rm w}\perp}^{\ \ i}\nabla_i\varrho\,. \tag{60}$$

It follows that, after a finite time interval during which the vortices have undergone a small cylindrical radial displacement, $\Delta\varrho$ say, the ensuing (local) change in the angular frequency lag will be given in terms of the corresponding (uniform) change $\Delta\Omega$ in the observable rotation frequency, namely that of the solid constituent $\Omega=\Omega_{\rm c}$ by the remarkably simple formula

$$\frac{\Delta[\Omega]}{\Omega}=-\frac{m}{m_\star}\left(\frac{\Delta\Omega}{\Omega}+2\frac{\Delta\varrho}{\varrho}\right)\,. \tag{61}$$

In the idealised case of pinning to an absolutely rigid solid the displacement $\Delta\varrho$ would of course vanish, so that $\Delta[\Omega]$ would be approximately determined as a function just of the spherical radial coordinate r by a proportionality relation of the form $\Delta[\Omega]\propto m/m_\star$, but in practice, even for perfect pinning, there will be deviations from this as the crustal rigidity modulus is expected [42] to be rather low (the solid structure will be effectively "squashy"). On the other hand if it is supposed that pinning is entirely ineffective (since the vortices may drift by either thermal [3] or quantum fluctuations [32]) then the generalised Proudman theorem derived below in Subsection 4.2 tels us that the angular velocity difference $[\Omega]$ – and hence also its variation $\Delta[\Omega]$ – will be a function not of r but of the cylindrical coordinate $\varrho = r\cos\theta$, so in that case (to allow for non-uniformity of $m_\star$) the contribution from the displacement term in (61) would have to be relatively significant.

3.2 Simplified global model

For the purpose of drawing conclusions from available empirical data such as that of Lyne et al. [33], rather than with the locally well defined but observationally inaccessible quantities involved in a formula such as (61) it can be more instructive in practice to work with corresponding crudely defined averages, using a total angular momentum decomposition of the form

$$J = J^{\rm f} + J^{\rm c} \tag{62}$$

in terms of "free" and "confined" contributions of the form

$$J^{\rm x} = \int n_{\rm x}\,\mu^{\rm x}_{\,i}\,\varrho^i\,{\rm d}^3 r\,, \tag{63}$$

with the "free" bayon momentum $\mu^{\rm f}_i$ given by (31) and with the "confined" baryon momentum given by tha analogous formula

$$\mu^{{\rm c}\,i} = m v_{\rm c}^{\,i} - m_\star^{\rm c}\,[v^i]\,, \qquad , \tag{64}$$

in which the effective mass function $m_\star^{\rm c}$ for the "confined" baryons is determined by its "free" baryon analogue,analogue $m_\star^{\rm f} = m_\star$, via the symmetric relation

$$n_{\rm c}(m_\star^{\rm c} - m) = n_{\rm f}(m_\star - m)\,. \tag{65}$$

Using an approximation in which $\Omega_{\rm f}$ is taken to be roughly uniform, as $\Omega_{\rm c}(=\Omega)$ must be in any case, these contributions will be given roughly by

$$J^{\rm f} \simeq I^{\rm ff}\Omega_{\rm f} + I^{\rm cf}\Omega_{\rm c}\,, \qquad J^{\rm c} \simeq I^{\rm cc}\Omega_{\rm c} + I^{\rm cf}\Omega_{\rm f}\,, \tag{66}$$

in terms of coefficients that combine to give the respective moments of inertia of the confined and free parts in the form

$$I^{\rm f} = I^{\rm ff} + I^{\rm cf}\,, \qquad I^{\rm c} = I^{\rm cc} + I^{\rm cf}\,. \tag{67}$$

In accordance with (64) the separate coefficients will be given, in terms of the cylindrical radius h, as the volume integrals

$$I^{\rm cc} = \int m_\star^{\rm c}\, n_{\rm c}\,\varrho^2\,{\rm d}^3 r\,, \qquad I^{\rm ff} = \int m_\star\, n_{\rm f}\,\varrho^2\,{\rm d}^3 r\,, \tag{68}$$

while, in view of the equivalence (65), the entrainment coefficient will be given by

$$I^{\rm cf} = \int (m - m_\star^{\rm c})\, n_{\rm c}\,\varrho^2\,{\rm d}^3 r = \int (m - m_\star)\, n_{\rm f}\,\varrho^2\,{\rm d}^3 r\,, \tag{69}$$

so the subtotals will be given by expressions of the familiar form

$$I^{\rm c} = m\int n_{\rm c}\,\varrho^2\,{\rm d}^3 r\,, \qquad I^{\rm f} = m\int n_{\rm f}\,\varrho^2\,{\rm d}^3 r\,, \tag{70}$$

which combine to give the complete stellar moment of inertia as

$$I = I^{\rm c} + I^{\rm f} = \int \rho\,\varrho^2\,{\rm d}^3 r\,. \tag{71}$$

In a simplified treatment it may be supposed that these moment of inertia coefficients remain constant (i.e. that the effects of a conceivable radial displacement $\Delta\,\varrho$ in (61) can be ignored) and that the superfluid contribution to the angular momentum is conserved during the interglitch period, $\Delta J^{\rm f}\simeq 0$, which by (66) evidently implies

$$\Delta\Omega_{\rm f} \simeq -\frac{I^{\rm cf}}{I^{\rm ff}}\,\Delta\Omega\,. \tag{72}$$

It may also be supposed that the total angular momentum variation is negligible, $\delta J = 0$ during the very short time interval in which the star undergoes a discontinuous glitch transformation involving an angular momentum transfer given by $\delta J^{\rm f}\simeq -\delta J^{\rm c}$ so that the corresponding angular velocity discontinuities will be related by

$$\delta\Omega_{\rm f} \simeq -\frac{I^{\rm c}}{I^{\rm f}}\,\delta\Omega\,. \tag{73}$$

To avoid a long term build up of too large a deviation of the superfluid angular velocity $\Omega_{\rm f}$ from the externally observable value, $\Omega_{\rm c} = \Omega$, the averages over many glitches of their total (continuous plus discontinuous) change per glitch should be the same,

$$\langle \Delta\Omega_{\rm f} + \delta\Omega_{\rm f}\rangle \simeq \langle \Delta\Omega + \delta\Omega\rangle\,, \tag{74}$$

a condition that is equivalently expessible using the notation of (59) as

$$\langle \delta[\Omega]\rangle \simeq -\langle \Delta[\Omega]\rangle\,. \tag{75}$$

The immediately preceding relations can be used to rewrite this condition in the form

$$\frac{\langle\delta\Omega\rangle}{\langle\Delta\Omega\rangle} \simeq -\frac{(I^{\rm f})^2}{I^{\rm ff}\,I}\,, \tag{76}$$

in which the left hand side contains only quantities that are directly observable, with values that have been typically found [33] to be given in order of magnitude by $\langle\delta\Omega\rangle/\langle\Delta\Omega\rangle \approx -10^{-2}$ in the case of Vela type pulsars. This magnitude has commonly been used as a basis for estimating the ratio $I^{\rm f}/I$ of the moment of inertia of the superfluid component involved to that of the whole star, on the basis of the usual supposition that there is no need to distinguish between the coefficients $I^{\rm f}$ and $I^{\rm ff}$. However it can be seen from (68) that the relation between these coefficients will actually be expressible in terms of a suitably weighted mean value $\bar m_\star$ of $m_\star$ by the formula

$$I^{\rm ff} \simeq \frac{\bar m_\star}{m} I^{\rm f}\,, \tag{77}$$

in which, according to the recent work [17] referred to above, the ratio $\bar m_\star/m$ is likely to be substantially larger than unity, which leads to a correspondingly augmented estimate,

$$\frac{I^{\rm f}}{I} \simeq -\frac{\bar m_\star}{m}\frac{\langle\delta\Omega\rangle}{\langle\Delta\Omega\rangle}\,, \tag{78}$$

for the relative value of the moment of inertia $I^{\rm f}$ of the relevant superfluid component as compared with the total moment of inertia I of the neutron star.

4 Stationary circularly symmetric configurations.

4.1 Total stress force in the general case

We now investigate the equilibrium conditions that need to be satisfied in the short and medium term when we neglect the slow secular evolution effects considered in the preceding section and consider the system to to be in a state that is exactly stationary, and with flow that is strictly circular in the sense that convective velocity contributions $v_{{\rm x}\perp}^{\,i}$ in (54) are absent, so that we simply have

$$v_{\rm f}^{\,i} = \Omega_{\rm f}\,\varrho^{\,i}\,, \qquad v_{\rm c}^{\,i} = \Omega_{\rm f}\,\varrho^{\,i}\,. \tag{79}$$

For a single-fluid model in such a non convective state the well known Proudman theorem tells us that the angular velocity must be uniform over each "Taylor cylinder" (as characterised by a fixed value of ϱ) for the case of a perfect fluid that is barotropic, meaning that there is no stratification. It will be shown in the present section how this result can be generalised to the two-fluid context under consideration here.

Using the circularity conditions (79) in conjunction with the axisymmetry conditions exemplified by (50), and assuming that the solidity property of the crust will ensure that the "normal" flow velocity of the "confined" current in such a stationary configuration will satisfy the rigidity condition, meaning that $\Omega_{\rm c}$ is uniform with the externally observed angular velocity Ω,

$$\nabla_i\,\Omega_{\rm c} = 0\,, \qquad \Omega_{\rm c} = \Omega\,, \tag{80}$$

it can be seen that the space parts of the force density formulae (24) and (26) will be expressible simply as

$$f^{\rm f}_{\,i} = n_{\rm f}\left(\nabla_i\,C_{\rm f} + \ell\,\nabla_i\,\Omega_{\rm f}\right)\,, \tag{81}$$

$$f^{\rm I}_{\,i} = n_{\rm c}\nabla_i\,C_{\rm c} + n_{\rm I}\nabla_i\,\tilde\mu^{\rm I}\,, \tag{82}$$

using the notation of Prix et al. [39] for the comoving particle energies, which are concisely specifiable for the respective "free" and "confined" constituents as the corresponding diagonal components

$$C_{\rm f} = \mathcal{E}^{\rm f}{}_{\rm f}\,, \qquad C_{\rm c} = \mathcal{E}^{\rm c}{}_{\rm c}\,, \tag{83}$$

of the (non-symmetric) energy matrix defined by the formula

$$\mathcal{E}^{\rm X}{}_{\rm Y} = -\pi^{\rm X}_{\,\nu}u^{\nu}_{\rm Y}\,. \tag{84}$$

This is another example of a formula that becomes more complicated and less heuristically meaningful when instead of using 4 dimensional notation it is written out using the traditional 3 dimensional notation, in which for example the comoving particle energy for the "free" constituent will be given by the formula

$$C_{\rm f} = m\phi + \tilde\chi^{\rm f} - \tfrac{1}{2}m\,v_{\rm f}^{\,2}\,, \tag{85}$$

in which, by (10) and (14), the relevant chemical potential will be given by

$$\tilde\chi^{\rm f} = \partial\tilde U_{\rm ins}/\partial n_{\rm f} - \tfrac{1}{2}[v]^2\partial\bar\rho_{\rm fc}/\partial n_{\rm f}\,, \tag{86}$$

with $\partial\bar\rho_{\rm fc}/\partial n_{\rm f} = n_{\rm f}\,\partial m_\star/\partial n_{\rm f} + m_\star - m$.

Without going into such details, it can be seen that the gravitational contributions to these comoving particle energies will cancel out in their difference, which will be given in terms of the chemical affinity (38) and the angular velocity difference $[\Omega] = \Omega_{\rm f} - \Omega$ by

$$C_{\rm c} - C_{\rm f} = \tilde{\mathcal{A}} + \ell\,[\Omega]\,. \tag{87}$$

If the crust were purely fluid, the force density $f^{\rm c}_{\,i}$ would just have to be equal in magnitude and opposite in sign to the force density acting on the superfluid, namely $-f^{\rm f}_{\,i}$. Detailed evaluation of the effects of solidity would require the use of a more elaborate model incorporating elastic rigidity, but as in the preceding work by Carter et al. [16] we can allow for the effect of finite rigidity by introducing a non vanishing stress force density f_i representing the discrepancy according to the definition (23), which can be rewritten in the form

$$\frac{1}{n_{\rm c}}\left(f_i - \frac{\rho}{\rho_{\rm f}} f^{\rm f}_{\,i}\right) = \frac{f^{\rm I}_{\,i}}{n_{\rm c}} - \frac{f^{\rm f}_{\,i}}{n_{\rm f}}\,, \tag{88}$$

in order to obtain a difference on the right hand side that is particularly amenable to simplification using (87). In terms of the chemical disequilibrium force density term defined by

$$f^{\chi}_{\,i} = n_{\rm c}\nabla_i\,\tilde{\mathcal{A}} + n_{\rm I}\nabla_i\tilde{\mu}^{\rm I}\,, \tag{89}$$

it can be seen that the formulae (81) and (82) lead to an expression of the remarkably simple form

$$\frac{f^{\rm I}_{\,i}}{n_{\rm c}} - \frac{f^{\rm f}_{\,i}}{n_{\rm f}} = \frac{f^{\chi}_{\,i}}{n_{\rm c}} + [\Omega]\nabla_i\ell\,. \tag{90}$$

We thus obtain an expression of the same form

$$f_i = \frac{\rho}{\rho_{\rm f}} f^{\rm f}_{\,i} + f^{\chi}_{\,i} + f^{\rm b}_{\,i}\,, \tag{91}$$

as that previously derived by Carter et al. [16] in a simplified treatment in which the effects of entrainment and stratification were not included, with the force density attributable to centrifugal buoyancy deficit given by the same formula as previously, namely

$$f^{\rm b}_{\,i} = n_{\rm c}[\Omega]\nabla_i\ell\,. \tag{92}$$

4.2 Generalised Proudman theorem for force-free superfluid case

Leaving aside the pinned case for treatment in the next subsection, let us restrict our attention in the remainder of the present subsection to the case in which the neutron superfluid flow is postulated to be force-free, which means that we have

$$f^{\rm f}_{\,i} = 0\,. \tag{93}$$

In this case the chemical and buoyancy contributions on the right of (91) will be the only sources for the stress force, which in this case will be given simply by

$$f_i = f^{\chi}_{\,i} + f^{\rm b}_{\,i}\,. \tag{94}$$

Under these circumstances (81) reduces to the form

$$\nabla_i\,C_{\rm f} + \ell\,\nabla_i\,[\Omega] = 0\,, \tag{95}$$

in which by (55) we have

$$\ell = \varrho^2\Big(m\,\Omega + m_\star[\Omega]\Big)\,. \tag{96}$$

Taking the exterior derivative of equation (95) implies that $\varepsilon^{ijk}(\nabla_i\ell)\nabla_j\,[\Omega]$ must vanish so that $\nabla_i\,[\Omega]$ has to lie in the direction of $\nabla_i\ell$ which is approximately that of $\nabla_i\varrho$. Consequently the condition (95) can be seen to provide us with a Proudman type theorem telling us that the energy parameter $C_{\rm f}$ and the angular velocity difference $[\Omega]$ must both be uniform over each of the deformed Tayor cylinders characterised by a fixed value of the angular momentum per "free" particle, ℓ. Provided the difference $[\Omega]$ is small compared with the uniform crust angular velocity Ω it can be seen from (96) that these modified Taylor cylinders will differ only by a small deformation from the ordinary Taylor cylinders that are characterised by fixed values of the cylindrical radial coordinate ϱ.

It is interesting to consider what would happen if the force given by (82) were also supposed to vanish, so that there would be no net force at all:

$$f^{\rm c}_{\,i} = 0 \Rightarrow A_{\rm c}\nabla_i\,C_{\rm c} + \nabla_i\,\tilde{\mu}^{\rm I} = 0\,, \tag{97}$$

where $A_{\rm c}$ is the nuclear mass number as introduced in (4). This would provide an analogue of the Proudman theorem, to the effect that the confined particle energy parameter $C_{\rm c}$ and the nuclear potential function $\tilde{\mu}^{\rm I}$ should both be uniform over surfaces characterised by a fixed value of $A_{\rm c}$.

A noteworthy special case of the application of the foregoing theorems is to the scenario considered by Prix et al [39] in which there is no stratification, meaning that $\nabla_i\,\tilde{\mu}^{\rm I} = 0$ and for which the differential rotation is also postulated to be uniform so that one has $\nabla_i\,[\Omega] = 0$, in which case it follows that in the absence of any (pinning or external stress) force both $C_{\rm c}$ and $C_{\rm f}$ would be uniform, as first integral constants of the motion, in the sense that their gradients too would vanish, $\nabla_i\,C_{\rm c} = \nabla_i\,C_{\rm f} = 0$.

Assuming that (as well as the gravitational background ϕ that may have been provided by a Cowling type approximation) the distributions of the angular velocities (or the corresponding angular momenta) and in our case also of the nuclear number density $n_{\rm I}$ (or the corresponding stratification potential $\tilde{\mu}^{\rm I}$) have been specified in advance to have values resulting from the previous history of the system, then the specification of two more independent quantities such as $C_{\rm c}$ and $C_{\rm f}$ will suffice to determine the system completely, as remarked by Prix et al. [39], who observed that their values as first integral constants could be fixed by the specification of the globally integrated totals of free and confined baryons.

A force free configuration with separate conservation of the free and confined constituents is not however what is most realistic in a system that is stationary over a secular timescale. The assumption of separate conservation of neutrons and protons is appropriate in the context of high frequency dynamical processes [18, 5] such as may give rise to a two stream instability [6]. However in an effectively stationary context it is to be expected that transfusion (by beta processes) would be able to establish the chemical equilibrium condition (44) to the effect that the affinity $\tilde{\mathcal{A}}$ should vanish, which by (87) means that one will have

$$C_{\rm c} = C_{\rm f} + \ell\,[\Omega]\,. \qquad (98)$$

In the special case for which $[\Omega]$ and hence (by the generalised Proudman theorem) also $C_{\rm f}$ are uniform it follows, since the distribution of ℓ will be highly non-uniform (roughly proportional to the squared cylindrical radius ϱ^2) that $C_{\rm c}$ will also be highly non-uniform. We thus obtain a scenario in which unlike that considered by Prix et al.[39] requires only a single constant of integration. This can be taken to be the globally integrated total baryon number, which suffices to determine the uniform value of $C_{\rm f}$, from which $C_{\rm c}$ will be obtainable unambiguously (without any need to specify any other arbitrary parameter) as the non-uniform field provided by substituting the relevant fixed values of $[\Omega]$ and $C_{\rm f}$. A scenario of this mathematically well defined type differs from the kind envisaged by Prix et al. [39] in that, despite the fact that no vortex pinning is involved, it requires that the solid structure of the crust should provide a stress force density that according to (94) will be given as a sum of stratification and buoyancy deficit forces by

$$f_i = n_{\rm I}\nabla_i\tilde{\mu}^{\rm I} + n_{\rm c}[\Omega]\nabla_i\ell\,. \qquad (99)$$

This could only vanish if $A_{\rm c}^{-1}\nabla_i\tilde{\mu}^{\rm I}$ were equal in magnitude and opposite in sign to $[\Omega]\nabla_i\ell$ which is mathematically conceivable but physically implausible (except near the stellar equator) because ℓ will depend mainly on the cylindrical coordinate ϱ, whereas one would expect that the other quantities involved would depend mainly on the spherical radial coordinate r.

4.3 Constants of integration for pinned superfluid case

To deal with the case when the superfluid vortices are pinned to the "normal" constituent it is convenient to focus attention not so much on the comoving diagonal element $C_{\rm f}$ in the energy matrix (84) but on the cross term

$$C = \mathcal{E}^{\rm f}{}_{\rm c} = C_{\rm f} + [v^i]\,\mu^{\rm f}{}_i\,, \qquad (100)$$

which is interpretable as the energy per particle of the "free" (superfluid) neutrons with respect to the"normal" rest frame of the "confined" baryons. Unlike the diagonal elements $C_{\rm f} = -\pi^{\rm f}{}_\nu u_{\rm f}^\nu$ and $C_{\rm c} = -\pi^{\rm c}{}_\nu u_{\rm c}^\nu$ (and the other off-diagonal element) this particular off-diagonal element $C = -\pi^{\rm f}{}_\nu u_{\rm c}^\nu$ has the property of being invariant with respect to the changes of chemical basis that were discussed in Subsection 2.6.

The usual supposition that, instead of moving freely, as was supposed in the preceding subsection, the vortices will satisfy the pinning condition expressed by (42) and (43) can be seen to be equivalent to the requirement that the "free" neutron current should be subject to a force density of the form

$$f^{\rm f}{}_i = -f^{\rm J}{}_i \qquad (101)$$

where $f^{\rm J}{}_i$ is the Magnus force density reacting on the solid structure of the confined constituent, as given in accordance with the usual Joukowsky formula (see appendix) by

$$f^{\rm J}{}_i = n_{\rm f}\varpi^{\rm f}{}_{ij}[v^j]\ , \qquad (102)$$

which works out in this stationary circularly symmetric case as

$$f^{\rm J}{}_i = n_{\rm f}[\Omega]\nabla_i\ell\,. \qquad (103)$$

It is to be noticed that, unlike the buoyancy deficit force density (92) to which it is proportional, this Magnus force density is invariant with respect to changes of chemical basis. When substituted in the general formula (91) for the total force acting on the system it provides an expression of the simple form

$$f{}_i = f^{\chi}{}_i - f^{\rm J}{}_i\,. \qquad (104)$$

By substituting the Joukowsky formula in (81) it can be seen that in terms of the crossed particle energy (100) the vortex pinning condition can be rewritten simply as

$$\nabla_i C = 0\,, \qquad (105)$$

which means that for any pinned configuration C will be a uniform integral constant, whose value can be fixed by the total globally integrated baryon number. In terms of this constant C the comoving energy per particle of the "free" constituent will be obtainable from (100) as the variable function of position given by

$$C_{\rm f} = C - [\Omega]\,\ell\,. \qquad (106)$$

It can be seen from (98) that the tranfusive equilibrium condition (44) will be expressible simply as

$$C_{\rm c} = C\,, \qquad (107)$$

which means that, when the specification of the system is completed by the imposition of this chemical equilibrium condition, then $C_{\rm c}$ will also have a uniform constant value.

The constancy of these quantities is interpretable as a modified Bernouilli theorem of a kind that is rather robust: it is to be emphasised that whereas the constancy deduced by Prix et al. [39] for $C_{\rm c}$ and $C_{\rm c}$ in the unpinned case was dependent on very restrictive assumptions about the uniformity of $[\Omega]$ and the negligibility of stratification, no such restrictions are involved in the derivation of constancy for $C_{\rm f}$ and $C_{\rm c}$ in the pinned case.

In this pinned equilibrium case, the analogue of the formula (99) for the stress force density that needs to be provided by the solidity of the crust will be given, as a difference between stratification and Magnus-Joukowsky contributions by the expression

$$f_i = n_{\rm I}\nabla_i\tilde\mu^{\rm I} - n_{\rm f}[\Omega]\nabla_i\ell\,, \tag{108}$$

which, unlike (99), has the noteworthy property of being chemically invariant.

The work in this section has used no approximations apart from the postulates of exact stationarity and axisymmetry, but if we make the realistic supposition that the configurations under consideration differ only slightly from a rigidly rotating reference state with fixed angular velocity $\bar\Omega$ then, to lowest order, we shall obtain the expressions

$$\ell = m\,\bar\Omega\,\varrho^2\,, \qquad\qquad \nabla_i\ell = 2m\,\bar\Omega\,\varrho\nabla_i\varrho\,. \tag{109}$$

This dependence of the angular momentum (per superfluid particle) ℓ on the cylindrical radial coodinate ϱ contrasts with the dependence mainly on the spherical radial coordinate r that one would expect for the number densities $n_{\rm I}$ and $n_{\rm f}$, as well as for the angular velocity difference $[\Omega]$ (which, in view of (61), might be expected to be roughly proportional to the depth dependent ratio $m/m_\star$). This is why, as in the case of (99), it is physically implausible that the gradient of the stratification potential $\tilde\mu^{\rm I}$ could be adjusted so as to cancel the terms in (108) except near the stellar equator.

5 Conclusions

The foregoing work shows how it is possible to construct an extensive range of simple, explicitly integrable, axisymmetric stratified two-fluid neutron star models in which a small differential angular velocity $[\Omega]$ can be chosen in advance as an arbitrary (implicitly history dependent) function of the spherical coordinates r and θ in the pinned case, and as an arbitrary function just of the cylindrical coordinate $\varrho = r\cos\theta$ in the unpinned case to which the generalised Proudman theorem applies.

This idealised category of non-dissipative models includes, as extreme case, the kind of force-free model constructed by Prix et al. [39] on the basis of the postulate of separate conservation of protons and neutrons. Except in this special case it is shown that the equilibrium of the two-fluid model necessitates the presence of a balancing force (given by (99) for unpinned configurations and by (108) for pinned configurations) that must presumably be provided by stress in the solid structure of the crust (whose detailed treatment is beyond the scope of the purely fluid description used here) which can build up only to a certain point at which the breakdown responsible for the glitch phenomenon will occur.

The exceptional case in which the presence of such (glitch producing) stress was avoided [39] depended on the postulate of separate conservation of protons and neutrons, a condition that would be realistic in the short timescale context of dynamic perturbations but not in the long timescale context of stationary configurations, for which the postulate of chemical equilibrium is advocated here as a more plausible idealisation.

It is to be understood that the force density given either by (99) or more likely by (108) will have to be balanced by a solid stress gradient, meaning that it will satisfy a relation of the form $f_i = \nabla_j\langle S\rangle^j{}_i$ in which $\langle S\rangle^j{}_i$ is a shear stress tensor that is the trace-free part of an elastic stress tensor $S^j{}_i$ of the kind recently described in the framework of an elastic solid model that is already available [10] for the description of the outer crust, of which an appropriate neutron conducting generalisation will soon be ready [9] for application to the inner crust (with density above the neutron drip threshold) under consideration here. The idea is that this shear stress tensor $\langle S\rangle_{ij}$ will be proportional to a trace-free shear strain tensor that can not excede a critical magnitude beyond which the solid structure will breakdown, and that a glitch will occur either when this critical value is reached or else, in the pinned case (108), at what may be an earlier stage when the Joukowski-Magnus force density (103) reaches a vortex slippage limit at which unpinning occurs.

For the study of processes occurring on intermediate timescales it will of course be necessary to include allowance for the dissipative processes that have been neglected in the idealised models considered here. As well as the controversial question of vortex slippage, [27, 26, 20, 21, 22] a particularly important issue concerns the timescales [41, 47, 45] of the weak interaction processes leading to chemical equilibrium.

Acknowledgments

The authors are grateful to Silvano Bonazzola, Loic Villain and Pawel Haensel for discussions concerning stratification and deviations from chemical equilibrium in neutron stars. Nicolas Chamel acknowledges support from the Lavoisier program of the French Ministry of Foreign Affairs.

References

[1] M. Alford, J. Bowers, K. Rajagopal, "Crystalline color superconductivity", *Phys. Rev. D***63** (2001), 074016.

[2] M.A. Alpar, H.F. Chau, K.S. Cheng, D. Pines, "Post-glitch relaxation of the VELA pulsar after its first eight large glitches - A reevaluation with the vortex creep model", *Ap. J.* **409** (1993) 345-359.

[3] M.A. lpar, D. Pines, P. .W Anderson, J. Shaham, "Vortex creep and the internal temperature of neutron stars. I - General theory", *Ap. J.***276** (1984) 325-334.

[4] M.A. Alpar, J.A. Sauls, "On the dynamical coupling between the superfluid interior and the crust of a neutron star" *Ap. J.* **327** (1988) 723-725.

[5] N. Andersson, G.L. Comer, D. Langlois, "Oscillations of general relativistic superfluid neutron stars", *Phys. Rev.* **D66** (2002) 104002.

[6] N. Andersson, G.L. Comer, R.Prix, "The superfluid two-stream instability and pulsar glitches", *M.N.R.A.S.* **354** (2004) 101-110.

[7] A. F. Andreev, E. P. Bashkin, "Three velocity hydrodynamics of superfluid solutions", *Sov. Phys. J.E.T.P.* **42** (1976) 164-167.

[8] G. Baym, C. Pethick, D. Pines, M. Ruderman, "Spin up in neutron stars: the future of the Vela pulsar", *Nature* **224** (1969) 872.

[9] B. Carter, E. Chachoua, "Newtonian mechanics of neutron superfluid in elastic star crust", ArXiv preprint astro-ph/0601658.

[10] B. Carter, E. Chachoua, N. Chamel, "Covariant Newtonian and Relativistic dynamics of (magneto)-elastic solid model for neutron star crust." *Gen. Rel. Grav.* **38** (2006) 83-119.

[11] B. Carter, N. Chamel, "Covariant analysis of Newtonian multifluid models for neutron stars: I Milne-Cartan structure and variational formulation", *Int. J. Mod. Phys. D* **13** (2004) 291-325.

[12] B. Carter, N. Chamel, "Covariant analysis of Newtonian multifluid models for neutron stars: II Stress - energy tensors and virial theorems; III Transvective, viscous and superfluid drag dissipation", *Int. J. Mod. Phys. D* **14** (2005) 717-774.

[13] B. Carter, N. Chamel, P. Haensel, "Entrainment coefficient and effective mass for conduction neutrons in neutron star crust: Simple nuclear models", *Nucl. Phys. A* **748** (2005) 675-697.

[14] B. Carter, N. Chamel, P. Haensel, "Effect of BCS pairing on entrainment in neutron superfluid current in neutron star crust", *Nucl. Phys. A* **759** (2005) 441-464.

[15] B. Carter, N. Chamel, P. Haensel, "Entrainment coefficient and effective mass for conduction neutrons in neutron star crust: Macroscopic treatment", ArXiv preprint astro-ph/0408083.

[16] B. Carter, D. Sedrakian, D. Langlois, "Centrifugal buoyancy as a mechanism for neutron star glitches", *A&A.* **361** (2000) 795 - 802.

[17] N. Chamel, "Band structure effects for dripped neutrons in neutron star crust", *Nucl. Phys. A* **747** (2005) 109-128.

[18] G.L. Comer, N. Andersson, "On the dynamics of superfluid neutron star cores", *M.N.R.A.S.* **328** (2001) 1129-1143.

[19] G.L. Comer, R. Joint, "A relativistic mean field model for entrainment in general relativity", *Phys. Rev.* **D68** (2003) 023002.

[20] D. Dall'Osso, G.L. Israel, L. Stella, A. Possenti, E. Perozzi, "The Glitches of the Anomalous X-Ray Pulsar 1RXS J170849.0-400910", *ApJ.***599** (2003) 485-497.

[21] P. Donati, P.M. Pizzochero, "Is there Nuclear Pinning of Vortices in Superfluid Pulsars?", *Phys. Rev. Lett.* **90** (2003), 211101

[22] P. Donati, P. M. Pizzochero, "Fully consistent semi-classical treatment of vortex-nucleus interaction in rotating neutron stars", *Nucl. Phys. A* **742** (2004), 363-379.

[23] L. M. Franco, B. Link, R. I. Epstein, "Quaking neutron stars" *Ap. J.* **543** (2000) 987-994.

[24] G. Hobbs, A. G. Lyne, B. C. Joshi, M. Kramer, I.H. Stairs, F. Camilo, R. N. Manchester, N. D'Amico, A. Possenti, V. M. Kaspi, "A very large glitch in PSR J1806-2125", *M.N.R.A.S.* **333** (2002) L7-L10.

[25] K. Iida & K. Sato, "Spin-down of Neutron Stars and Compositional Transitions in the Cold Crustal Matter", *Ap. J.* **477** (1997) 294-312.

[26] D. I. Jones & N. Andersson, "Freely precessing neutron stars: model and observations", *M.N.R.A.S.* **324** (2001) 811-824.

[27] P. B. Jones, "The origin of pulsar glitches", *M.N.R.A.S.* **296** (1998) 217-224.

[28] V. M. Kaspi, J. R. Lackey, D. Chakrabarty, "A Glitch in an Anomalous X-Ray Pulsar", *Ap. J.* **537** (2000) L31-L34.

[29] V. M. Kaspi, F. P. Gavriil, "A Second Glitch from the "Anomalous" X-Ray Pulsar 1RXS J 170849.0-4000910", *Ap. J.* **596** (2003) L71-L74.

[30] D. Langlois, D. Sedrakian, B. Carter, "Differential rotation of relativistic superfluid in neutron stars", *MNRAS* **297** (1998) 1198-1201.

[31] B. Link, "Constraining Hadronic Superfluidity with Neutron Star Precession", *Phys. Rev. Lett.* **91** (2003) 101101.

[32] B. Link, R. I. Epstein, G. Baym, "Superfluid vortex creep and the rotational dynamics of neutron stars", *ApJ.***403** (1993) 285-302.

[33] A.G. Lyne, S. L. Shemar, F. G. Smith, "Statistical studies of pulsar glitches", *M.N.R.A.S.* **315** (2000) 534-542.

[34] G. Mendell, "Superfluid hydrodynamics in rotating neutron stars. I Nondissipative equations", *Ap. J.* **380** (1991) 515-529.

[35] M. Morii, N. Kawai, N. Shibazaki, "Pulse Profile Change Possibly Associated with a Glitch in an Anomalous X-Ray Pulsar 4U 0142+61", *Ap. J.* **622** (2005) 544-548.

[36] M.A. Perez-Garcia, N. Corte, L. Mornas, J.P. Suares-Curieses, J. Diaz-Alonso, "Formation of an ordered phase in nuclear matter" *Nucl. Phys. A* **699** (2002) 939.

[37] P.M. Pizzochero, L. Viverit, R.A. Broglia, "Vortex nucleon interaction and pinning forces in neutron stars" *Phys. Rev. Lett.* **79** (1997) 3347-3350.

[38] R. Prix, "Slowly rotating two-fluid neutron star model" *A&A.* **352** (2000) 623 - 631.

[39] R. Prix, G.L. Comer, N. Andersson, "Slowly rotating superfluid neutron star model with entrainment", *A&A.* **381** (2002) 178-196.

[40] R. Prix, J. Novak, G.L. Comer, "Relativistic numerical models for stationary superflid neutron stars" *Phys. Rev.* **D71** (2005) 043005.

[41] A. Reisenegger, "Constraining Dense Matter Superfluidity through Thermal Emission f *Ap. J.* **485** (1997) 313-318.

[42] M. Ruderman, "Neutron star crustal plate tectonics. I. Magnetic dipole evolution in lillisecond pulsars and low-mass X-ray binaries", *Ap. J.* **366** (1991) 261-269.

[43] M. Ruderman, T. Zhou, K. Chen, "Neutron star magnetic field evolution, crust movement and glitches", *Ap. J.* **492** (1998), 267-280.

[44] J.A. Sauls, in *Timing Neutron Stars* ed. H. Ogelman, E.D.P. Van den Heuvel (Kluwer, Dordrecht, 1989) 457-490.

[45] L. Villain & P. Haensel, "Non equilibrium beta processes in superfluid neutron star cores", ArXiv preprint astro-ph/0504572.

[46] N. Wang, R.N. Manchester, R.T. Pace, M. Bailes, V.M. Kaspi, B. W. Stappers, A. G. Lyne, "Glitches in southern pulsars", *M.N.R.A.S.* **317** (1998) 843-860.

[47] D. G. Yakovlev, A.D. Kaminker, O. Y. Gnedin, P. Haensel, "Neutrino emission from neutron stars", *Phys. Rep.* **354** (2001) 1-155.

6 Appendix

6.1 Joukowsky formula for Magnus force

The purpose of this appendix is to derive the formula used above for the force exerted on the fluid in the pinned case and to verify that it does indeed balance the collective effect of the Magnus forces exerted on the individual vortices. We shall start by evaluating the latter, using the generalised Joukowski formula that has recently been derived [12] for the force exerted on an isolated vortex by a mixture of *irrotational perfect* fluids.

The present application is to the case of a vortex that is fixed with respect to the crust rest frame characterised by the flow 4 vector $u_{\rm c}^{\mu}$ of the "normal" constituent, and that is oriented in the direction specified by a spacelike unit 4-vector ν^{μ} say, for which the force per unit length will be given by

$$\mathcal{F}_{\nu} = \mathcal{C}^{\rm f} n^{\mu} \varepsilon_{\mu\nu\rho\sigma} \nu^{\rho} u_{\rm c}^{\sigma} \tag{110}$$

where n^{μ} is the (chemical gauge invariant) relative flow current vector of the superfluid and $\mathcal{C}$ is its (chemical gauge invariant) circulation as given by

$$\mathcal{C}^{\rm f} = \oint \pi^{\rm f}{}_\mu {\rm d}x^\mu \,, \tag{111}$$

in which the integral is carried out along any *closed* path surrounding the vortex.

Since the effect of many such vortices in closely spaced parallel array will be represented by a vorticity vector

$$w^\mu = w\,\nu^\mu \tag{112}$$

of the type introduced in (40), whose magnitude w is the value of the total circulation per unit area, it follows that the corresponding force density (obtained as force per unit length per unit area) will be given by

$$f^{\rm J}{}_\nu = \frac{w}{\mathcal{C}^{\rm f}} \mathcal{F}_\nu \tag{113}$$

so we end up with the formula

$$f^{\rm J}{}_\nu = n^\mu \varepsilon_{\mu\nu\rho\sigma} w^\rho u_{\rm c}^\sigma \tag{114}$$

in which it is to be observed that the value of the circulation $\mathcal{C}^{\rm f}$ round an individual vortex has cancelled out. (Thus it is not necessary to know that it will actually be given in terms of the Dirac-Planck constant by the formula $\mathcal{C}^{\rm f} = \pi\hbar$ in which the usual factor 2 is missing because of the pairing due the fermionic nature of the neutrons.)

To obtain the force density on the fluid we start from the formula (24) in which, since we are considering a stationary axisymmetric situation, the divergence term will automatically drop out leaving the expression

$$f^{\rm f}{}_\mu = n_{\rm f}^\nu \varpi^{\rm f}{}_{\nu\mu} \,. \tag{115}$$

which would be equivalent just to the ordinary Euler equation if the force term vanished. It is evident that the pinning condition

$$u_{\rm c}^\nu \varpi^{\rm f}{}_{\nu\mu} = 0, \tag{116}$$

can be used to rewrite (115) in the form

$$f^{\rm f}{}_\mu = n_{\rm f}(u_{\rm f}^\nu - u_{\rm c}^\nu)\varpi^{\rm f}_{\nu\mu} = n^\nu \varpi^{\rm f}{}_{\nu\mu} \,. \tag{117}$$

It can also be seen to follow from (116) that, in terms of the vector introduced in (40), the vorticity form will be expressible as

$$\varpi^{\rm f}{}_{\nu\mu} = \varepsilon_{\mu\nu\rho\sigma} w^\rho u_{\rm c}^\sigma \,, \tag{118}$$

so that (114) can be rewritten in the form

$$f^{\rm J}{}_\mu = -n^\nu \varpi^{\rm f}{}_{\nu\mu} \,, \tag{119}$$

which evidently does indeed exactly balance (117).

6.2 Effect of entrainment on vortex density

It is of interest to consider the effect of entrainment on the vorticity magnitude w which is the circulation per unit area round an infinitesimal circuit orthogonal to the local axis of rotation. Remarking that the circulation $\mathcal{C}^{\rm f}$ is simply given by $2\pi\ell$, where ℓ is the angular momentum per free particle, it can be seen from (55), by considering the circulation between nearby circuits with crylindrical radius values ϱ and $\varrho + \delta\varrho$, that the local vorticity magnitude will be expressible in terms of the angular velocity difference $[\Omega] = \Omega_{\rm f} - \Omega_{\rm c}$ and the mass increment $m^{\rm f}_{\rm c} = m_\star - m$ as

$$w = \frac{\delta\mathcal{C}^{\rm f}}{2\pi\varrho\delta\varrho} = 2m\Omega_{\rm f} + 2m^{\rm f}_{\rm c}[\Omega] + \varrho\frac{\delta(m_\star[\Omega])}{\delta\varrho} \,. \tag{120}$$

Knowing that the circulation round an individual quantised vortex line in the fermionic superfluid condensate will be given by $\mathcal{C}^{\rm f} = \pi\hbar$ we see that the corresponding surface number density of quantised vortex lines will simply be given by $w/\mathcal{C}^{\rm f} = w/(\pi\hbar)$. The first term in equation (120) can be recognised as the well-known formula for a single superfluid.

In practice, the extra terms proportional to $[\Omega]$ and its gradient will be negligible for the astrophysical applications under consideration, in which we shall generally have $[\Omega] \ll \Omega$, provided we are using an "operational" gauge in which magnitude of the effective mass $m_\star$ will remain comparable with the ordinary baryonic mass m. If the evaluation were carried out in a "comprehensive" gauge, meaning a chemical basis in which all the neutrons are deemed to be "free", the relevant value of the effective mass would become extremely large near the "neutron drip" transition, but the value of the (gauge dependent) difference $[\Omega]$ would correspondingly become even smaller, since the product $m_\star[\Omega]$ is chemically invariant, so that the same value would ultimately be obtained for the vorticity magnitude w itself.